



Abiotic features

- low intertidal flats (<50% E.)
high intertidal flats (>50% E.)
- sl sh sand (<5% clay)
xl xh muddy sand (5-8% clay)
ml mh sand (>8% clay)
- 67% emersion time
- 0 beach above Mean High Tide
- N depth in m below Sea Level < 10
- N10 depth in m below Sea Level > 10

Biotic features

- musselbed
- Zostera marina bed (with 200 meter buffer)
- Zostera noltii (with 200 meter buffer)
- 1 mud- or sand flat with salt-marsh pioneers
- 2 low salt marsh (Puccinellion maritima)
- 3 middle salt marsh (Armerion maritima)
- 4 high marsh (Lolio-Potentillion anserinae)
- 5 transition between salt-marsh and sand dunes
- 6 reed marshes

**HABITATS OF THE NETHERLANDS,
GERMAN AND DANISH WADDEN SEA**

Compiled by: K. S. Dijkema, Rijksinstituut voor Natuurbeheer, Texel

1 : 100.000 sheet 4

ALTERA
WAGeningenUR

nature-consult

N
GIS-Compilation:
WGS_1984_UTM_Zone_32N
0 1 2 3 4 5
Kilometer

H.-P. Dauck
nature-consult
Hackelbrink 21
D-31139 Hildesheim
www.nature-consult.de